arXiv:1302.0013v1 [cond-mat.stat-mech] 31 Jan 2013

Ground-state Energies of Spinless Free Fermions and Hard-core Bosons

Wenxing Nie,[1] Hosho Katsura,[2] and Masaki Oshikawa[1]
[1] *Institute for Solid State Physics, University of Tokyo, Kashiwa 277-8581, Japan*
[2] *Department of Physics, Gakushuin University, Tokyo 171-8588, Japan*

(Dated: February 4, 2013)

We compare the groundstate energies of bosons and fermions with the same form of the Hamiltonian. If both are noninteracting, the groundstate energy of bosons is always lower, owing to Bose-Einstein Condensation. However, the comparison is nontrivial when bosons do interact. We first prove that, when the hopping is unfrustrated (all the hopping amplitudes are non-negative), hard-core bosons still must have a lower groundstate energy than the fermions. When the hopping is frustrated, bosons can have higher groundstate energy than fermions. We prove rigorously that this inversion indeed occurs in several examples, using various techniques.

Introduction. — Statistics of identical particles is one of the most fundamental concepts in quantum physics. One of the consequences of the particle statistics appears in the groundstate energy. For a system of free particles, the groundstate of bosons is obtained by putting all the particles in the lowest-energy state of the single-particle Hamiltonian, while the groundstate of fermions is obtained by filling the individual single-particle states up to the Fermi level (Pauli exclusion principle). Thus, the groundstate energy of bosons E_0^{B} and that of fermions E_0^{F}, for the same form of the Hamiltonian, satisfy

$$E_0^{\mathrm{B}} \le E_0^{\mathrm{F}}, \tag{1}$$

if the particles are non-interacting.

The comparison becomes nontrivial when the particles do interact; the Bose-Einstein condensation is no longer perfect in interacting systems. Intuitively, it would be still natural to expect that Eq. (1) holds. However, recently an apparent counterexample was found numerically [1, 2]. This motivates us to examine the fundamental question how general the "natural" inequality (1) is and when it can be actually violated.

To simplify the matter, in this paper we focus on the comparison of hard-core bosons with spinless fermions. The Hamiltonian is given as

$$\mathcal{H} = -\sum_{j,k}\left(t_{jk}c_j^\dagger c_k + \text{H.c.}\right) - \sum_j \mu_j n_j + \sum_{j,k} V_{jk} n_j n_k, \tag{2}$$

where each site j belongs to a lattice Λ, $n_j \equiv c_j^\dagger c_j$, and $t_{jk} = 0$ is assumed for $j = k$. The uniform (site independent) part of μ_j is the chemical potential μ. For a system of fermions, we identify c_j with the fermion annihilation operator f_j satisfying the standard anticommutation relations. For a system of bosons, we identify c_j with the boson annihilation operator b_j satisfying the standard commutation relations, with the hard-core constraint $n_j = 0, 1$ at each site. The hard-core constraint may be implemented by introducing the on-site interaction $U\sum_j n_j(n_j - 1)$ and then taking $U \to \infty$.

We note that, the Hamiltonian (2) conserves the total particle number. Thus the groundstate can be defined for a given number of particles M (canonical ensemble), or for a given chemical potential μ (grand-canonical ensemble). The comparison between bosons and fermions can be made in either circumstance.

Natural Inequality. — First we present a sufficient condition for the "natural" inequality (1) to hold. (See Ref. [3] for a similar inequality for spinful fermions).

Theorem 1. *The inequality* (1) *holds for any given number of particles M on a finite lattice Λ, if all the hopping amplitudes t_{jk} are real and non-negative.*

Proof. Let us take the occupation number basis $|\phi^a\rangle \equiv |\{n_j^a\}\rangle$, where $\sum_j n_j^a = M$. The number operator n_j has the same matrix element in this basis, for hard-core bosons and spinless fermions. On the other hand, for bosons, the matrix elements of each hopping term in the Hamiltonian (2) is non-positive in this basis. Thus matrix elements for bosons $\mathcal{H}^{\mathrm{B}}_{ab}$ and those for fermions $\mathcal{H}^{\mathrm{F}}_{ab}$ satisfy

$$\mathcal{H}^{\mathrm{B}}_{ab} = \begin{cases} -|\mathcal{H}^{\mathrm{F}}_{ab}| & (a \neq b) \\ \mathcal{H}^{\mathrm{F}}_{aa} & (a = b) \end{cases} \tag{3}$$

Let $|\Psi_0\rangle_{\mathrm{F}} = \sum_a \psi_a |\phi^a\rangle_{\mathrm{F}}$ be the (normalized) groundstate for fermions. To obtain an upper bound on the groundstate energy of bosons, we consider the trial state $|\Psi_0\rangle_{\mathrm{B}} = \sum_a |\psi_a||\phi^a\rangle_{\mathrm{B}}$, where $|\phi^a\rangle_{\mathrm{B}}$ is the basis state for bosons corresponding to $|\phi^a\rangle_{\mathrm{F}}$. Then Eq. (3) implies

$$E_0^{\mathrm{B}} \le {}_{\mathrm{B}}\langle\Psi_0|\mathcal{H}|\Psi_0\rangle_{\mathrm{B}} \le {}_{\mathrm{F}}\langle\Psi_0|\mathcal{H}|\Psi_0\rangle_{\mathrm{F}} = E_0^{\mathrm{F}}. \tag{4}$$

□

As a simple corollary, the groundstate energies for a given chemical potential μ also satisfy Eq. (1).

The sign of hopping amplitudes in a system of bosons is related to frustration. This is clear if one maps the system of hard-core bosons to the quantum spin system with $S = 1/2$; positive hopping amplitudes t_{jk} correspond to ferromagnetic XY interaction. Theorem 1 means that, if there is no frustration in the hopping term, hard-core bosons always have lower energy than corresponding fermions. This also gives an alternative understanding of why the fermions have higher energy in the "unfrustrated" case: statistical phases of fermions introduce a kind of frustration to the system.

Let us now discuss how the natural inequality (1) can be violated. According to Theorem 1, in order to realize the violation, it is necessary to introduce a frustration by choosing some amplitudes t_{jk} to be negative or complex. While the presence of frustration is not a sufficient condition, we will demonstrate that the violation indeed occurs in several concrete examples. In the following, for simplicity we only discuss tight-binding models of hard-core bosons and corresponding noninteracting fermions, setting $V_{jk} = 0$.

Particles on a Ring. — We begin with a simple but instructive example in one dimension: tight-binding model on a ring $\mathcal{H} = -\sum_{j=1}^{N} (c_j^\dagger c_{j+1} + \text{H.c.})$. The hard-core boson version of this model, which is equivalent to the $S = 1/2$ XY chain, can be mapped to the model of noninteracting fermions on the ring by the Jordan-Wigner transformation [4, 5]. Thus the hard-core bosons and fermions are almost equivalent in this case.

However, a care should be taken on the boundary condition when we discuss the ring of finite length. For simplicity, here we assume that the number of sites N is an integral multiple of 4, and the number of particles $M = N/2$ (an even integer by assumption). Then the hard-core bosons with the periodic (antiperiodic) boundary condition is mapped to noninteracting fermions with the antiperiodic (periodic, respectively) boundary conditions. Noninteracting fermions on a ring have a groundstate energy density lower by $O(1/N^2)$ for the antiperiodic boundary condition, compared to the periodic boundary condition [6]. This implies that the hard-core bosons have lower energy than fermions on a ring with the periodic boundary condition, conforming to Theorem 1 since all the hopping amplitudes are non-negative. On the other hand, the same result implies that, under the antiperiodic boundary condition, the hard-core bosons have higher energy than fermions. Imposing the antiperiodic boundary condition is equivalent to introducing a π-flux inside the ring, which can cancel the effect of the statistical phase so that the inequality (1) is indeed inverted. The energy difference on the ring, however, vanishes asymptotically in the thermodynamic limit $N \to \infty$. Thus, we shall seek for different examples where the hard-core bosons have higher energy than fermions in the thermodynamic limit.

Two Dimensional System with Flux. — A natural system to consider would be a two-dimensional lattice with flux. Under the periodic boundary condition, the total flux is quantized as integer flux quanta (the unit flux quantum is 2π in our unit). For the finite square lattice with N plaquettes, the flux per plaquette Φ is quantized in unit of $2\pi/N$. Such a uniform flux can be represented using the string gauge [7]. We obtained the groundstate energy of hard-core bosons and fermions with various densities of particles and various values of flux using exact numerical diagonalization, for square lattices with periodic boundary conditions, up to 26 sites. The result for the $\sqrt{26} \times \sqrt{26}$ square lattice is shown in Fig. 1.



Figure 1. Difference of groundstate energy density between the hard-core bosons and fermions ($\Delta\epsilon = E_0^{\text{B}}/N - E_0^{\text{F}}/N$) on the $\sqrt{26} \times \sqrt{26}$ square lattice with Φ flux per plaquette and n_e particle per site. The natural inequality (1) holds in white region, while its violation is color coded. Statistical transmutation is expected along the two solid lines $\Phi = n_e$ and $\Phi = 1 - n_e$.

We find that the "natural" inequality (1) is violated in a region of the phase diagram. In particular, the inversion is significant along the diagonal lines $\Phi = n_e$ and $\Phi = 1 - n_e$. These lines are precisely where the statistical transmutation between the hard-core boson and the fermion is expected to occur [8, 9]. Namely, in the mean-field level, one flux quantum can be attached to each particle, transforming fermions into bosons and vice versa, at the same time eliminating the background field. At zero field, the frustration is absent and hard-core bosons have lower energy than fermions. Thus, the statistical transmutation implies that, hard-core bosons have higher energy than fermions on the lines $\Phi = n_e$ and $\Phi = 1 - n_e$. While this argument is not rigorous, our numerical result supports the statistical transmutation scenario. (For a related discussion for spinful electrons, see Ref. [10].) Numerical results for the square lattices of various sizes up to 26 sites (not shown) suggest that the energy difference is nonvanishing in the thermodynamic limit. Furthermore, in the following, we rigorously prove the inversion of the groundstate energies for several lattices in the thermodynamic limit.

Flat Band. — Some models of noninteracting fermions have completely flat bands as the lowest energy band. In these models, as long as all the fermions occupy the lowest flat band, Pauli exclusion principle plays no role in increasing the groundstate energy. Thus, such flat band models would have better chance to realize the inversion of the groundstate energies. Here we show that the inequality (1) is indeed violated in a few examples with flat bands.

First we discuss the delta-chain model, for which the violation of Eq. (1) was numerically found for small clusters [1, 2]. The Hamiltonian of the model can be written in the following form [11, 12]:

$$\mathcal{H} = \sum_{j=1}^{N} a_j^\dagger a_j, \tag{5}$$

where the a-operator, which acts on each triangle, is defined as $a_j = c_{2j-1} + \sqrt{2}c_{2j} + c_{2j+1}$. Here we use a periodic boundary condition to identify c_{2N+1} with c_1. The Hamiltonian $\mathcal{H}$ corresponds to a model with negative hopping amplitudes t_{jk} (as defined in Eq. (2)), which lead to frustration.

The model in the single-particle sector has two bands. The lower flat band with zero energy is spanned by states annihilated by a_j's. We note that the Hamiltonian (5) is modified from that in Ref. [1] by a constant chemical potential, so that the flat band has exactly zero energy. Thus the groundstate energy of the fermionic version of the model (5) is zero as long as the filling fraction (particle number per site) ν satisfies $\nu \leq 1/2$. On the other hand, the groundstate energy E_0^{B} of bosons is zero as long as $\nu \leq 1/4$ since the localized zero-energy states do not overlap each other in this range of filling [13].

Now let us derive a nontrivial lower bound for E_0^{B} for filling fractions $\nu > 1/4$. We decompose the model into clusters, each containing p unit cells:

$$\mathcal{H} = \sum_{n=0}^{N/p-1} \mathcal{H}_n^{(p)} + \sum_{n=1}^{N/p} a_{np}^\dagger a_{np}, \tag{6}$$

where $\mathcal{H}_n^{(p)} = \sum_{j=1}^{p-1} a_{np+j}^\dagger a_{np+j}$ is the Hamiltonian for the $(n+1)$-th cluster. Since the second term $\sum_{n=1}^{N/p} a_{np}^\dagger a_{np}$, which connects neighboring clusters, is positive semidefinite, the groundstate energy $\tilde{E}_0^{\mathrm{B}}$ of the first term $\tilde{\mathcal{H}} = \sum_{n=0}^{N/p-1} \mathcal{H}_n^{(p)}$ satisfies $\tilde{E}_0^{\mathrm{B}} \leq E_0^{\mathrm{B}}$. $\tilde{\mathcal{H}}$ is a sum of mutually commuting cluster Hamiltonians $\mathcal{H}_n^{(p)}$. Thus $\tilde{E}_0^{\mathrm{B}}$ is simply given by the sum of groundstate energies of each cluster. The particle number within each cluster is also conserved separately in $\tilde{\mathcal{H}}$. Let us choose $p = 4$ as in Fig. 2. The cluster contains 8 sites and the number of particles is $m = 0, 1, \ldots, 8$. We obtained the groundstate energy $\epsilon_0^{(p)}(m)$ of the cluster in each particle number m sector, by numerical exact diagonalization. It is found that $\epsilon_0^{(4)}(m) = 0$ for $0 \leq m \leq 3$, and $\epsilon_0^{(4)}(m) \geq \Delta_{\mathrm{DC}}^{(4)} = 0.372605$ for $4 \leq m \leq 8$.

If we consider the filling fraction in the range $3/8 < \nu \leq 1/2$, it follows from Dirichlet's box principle that there is at least one cluster which contains 4 or more particles. Thus, in this range, $\tilde{E}_0^{\mathrm{B}} \geq \Delta_{\mathrm{DC}}^{(4)}$ for any system size N, while $E_0^{\mathrm{F}} = 0$. Therefore, the inversion of the groundstate energies holds also in the thermodynamic limit.



Figure 2. Decomposition of the delta-chain Hamiltonian to clusters. Each cluster contains $p = 4$ unit cells (8 sites), including one decoupled site at the top of the dashed triangle. The hoppings on dashed triangles $\sum_{n=1}^{N/p} a_{np}^\dagger a_{np}$ are omitted from the Hamiltonian. Since the omitted terms are positive semidefinite, the groundstate energy for the decoupled Hamiltonian $\tilde{\mathcal{H}}$ gives a lower bound for the groundstate energy of the original Hamiltonian $\mathcal{H}$.

This method can be easily extended to other lattices. For example, the standard nearest-neighbor hopping model on a kagome lattice is described by

$$\mathcal{H} = \sum_{\alpha} a_{\triangle_\alpha}^\dagger a_{\triangle_\alpha} + \sum_{\alpha} a_{\nabla_\alpha}^\dagger a_{\nabla_\alpha}, \tag{7}$$

where $\triangle_\alpha$ and ∇_α are elementary triangles pointing up and down respectively, of the kagome lattice. We define $a_{\triangle_\alpha} \equiv c_{\alpha_1} + c_{\alpha_2} + c_{\alpha_3}$, where $\alpha_{1,2,3}$ refer to the three sites belonging to $\triangle_\alpha$, and likewise for a_{∇_α}. The fermionic version of the model has three bands, the lowest of which is a flat band at zero energy [13–15]. Thus $E_0^{\mathrm{F}} = 0$ when $\nu \leq 1/3$. For the groundstate energy of the bosonic version, we can use the cluster decomposition technique similar to what we have discussed above for the delta-chain. Let us choose the 12-site cluster of the "Star of David" shape. The groundstate energy ϵ_0 of each cluster is zero in the fixed particle number sectors with $m \leq 3$, but is positive in the sectors with $m \geq 4$. Thus, invoking Dirichlet's box principle again, Eq. (1) is violated for filling fraction $1/4 < \nu \leq 1/3$.

The outcome of the above argument depends on the cluster size taken. In fact, the range of filling fraction ν for which we have proved the violation of Eq. (1) is not optimal. In an alternative approach generalizing the techniques used in the context of flat-band ferromagnetism [11] and of frustrated antiferromagnets near the saturation field [13], we can extend the region of ν where Eq. (1) is violated [16]. For example, in the delta-chain model, Eq. (1) is violated for $1/4 < \nu \leq 1/2$.

Anderson's argument. — In order to deal with more general models, we develop an alternative approach. Let us discuss the square lattice with π-flux per plaquette, at zero chemical potential. The Hamiltonian reads

$$\mathcal{H} = -\sum_{<j,k>} (t_{jk} c_j^\dagger c_k + \mathrm{H.c.}), \tag{8}$$

where $t_{jk} = \pm 1$ as specified in Fig. 3. We note that Lieb has shown that π-flux minimizes the groundstate energy of fermions on the square lattice [17]. On the other hand, an argument similar to the Proof of Theorem 1 can be used to prove a lattice version of Simon's theorem on diamagnetism of bosons [18]. Namely, for bosons, introduction of flux always increases the groundstate energy. These, together with the statistical transmutation argu-



Figure 3. The square lattice with π flux in each plaquette. We choose the gauge so that the hopping amplitude t_{jk} is $+1$ on black links, and -1 on blue ones. The brown cross represents a cluster of 12 sites.

ment discussed earlier, suggest a possibility of violation of Eq. (1) with π-flux per plaquette.

The fermionic version of this model does not have a flat band. Nevertheless, from the exact dispersion relation, the groundstate energy at half filling ($\mu = 0$) is obtained exactly as $E_0^{\mathrm{F}} \sim -0.958091N$ in the limit of large N. Now we turn to the "grandcanonical" groundstate energy of the corresponding boson model at the same chemical potential ($\mu = 0$). Here we use Anderson's argument [19–21] by writing the Hamiltonian as a sum of sub-Hamiltonians:

$$\mathcal{H} = \sum_{\alpha} h_{\alpha}. \tag{9}$$

The groundstate energy E_0 of $\mathcal{H}$ satisfies $E_0 \geq \sum_{\alpha} \epsilon_0^{\alpha}$, where ϵ_0^{α} is the groundstate energy of h_{α}. For the present problem, we can choose h_{α} as

$$h_{\alpha} = -\frac{1}{2} \sum_{\langle j,k \rangle \in +_{\alpha}} (t_{jk} c_j^{\dagger} c_k + \text{H.c.}), \tag{10}$$

where $+_{\alpha}$ refers to a cross-shaped cluster of 12 sites as shown in Fig. 3. We consider all the clusters with the same pattern of hopping amplitudes within the cluster, in the square lattice. As a consequence, each cluster overlaps with 4 other clusters and each link appears in two different clusters when periodic boundary conditions are imposed. The factor 1/2 in Eq. (10) compensates this double counting. The grandcanonical groundstate energy of the cross-shaped cluster is obtained by exact diagonalization as $\epsilon_0^{\alpha} = -3.609035$.

Since there are $N/4$ such clusters in the square lattice of N sites, we obtain

$$E_0^{\mathrm{B}}/N \geq -3.609035/4 = -0.902259 > E_0^{\mathrm{F}}/N. \tag{11}$$

Thus the inversion of the groundstate energies is proved for the square-lattice π-flux model with $\mu = 0$. This relation, expected from the statistical transmutation argument discussed earlier, is now proved rigorously.

This argument is not restricted to two dimensional systems. Let us consider the standard tight-binding model on a pyrochlore lattice: $\mathcal{H} = \sum_{\langle j,k \rangle} (c_j^{\dagger} c_k + \text{H.c.})$, where $\langle j,k \rangle$ runs over all pairs of nearest-neighbor sites in the three dimensional pyrochlore lattice. Again we set the chemical potential $\mu = 0$. We note that this model has frustrated hoppings with this choice of the sign. The model in the single-particle sector has two degenerate flat bands at the energy $\mathcal{E} = -2$ and two dispersive bands touching the flat bands [15]. Thus for fermions, the groundstate energy at $\mu = 0$ satisfies $E_0^{\mathrm{F}} < -2(N/2) = -N$, where N is the number of sites of the lattice.

The hard-core boson version of this model can be decomposed as Eq. (9) with $h_{\alpha} = (1/4) \sum_{\langle j,k \rangle \in \mathrm{TD}_{\alpha}} (c_j^{\dagger} c_k + \text{H.c.})$, where TD_{α} refers to each dimer of elementary tetrahedra of the pyrochlore lattice sharing a vertex (site). Here we count dimers in any direction; each tetrahedron (and thus each link) belongs to 4 dimers. The factor 1/4 in the definition of h_{α} is introduced to compensate the overcounting. The groundstate energy of a tetrahedra dimer is obtained by exact diagonalization as $\epsilon_0^{\alpha} = -(2+\sqrt{2})/4 = -0.853554$. Since there are N dimers of tetrahedra, the groundstate energy of bosons at $\mu = 0$ satisfies

$$E_0^{\mathrm{B}}/N \geq -(2+\sqrt{2})/4 > E_0^{\mathrm{F}}/N. \tag{12}$$

Thus we have proved the violation of Eq. (1) for the simple tight-binding model on the three-dimensional pyrochlore lattice.

Conclusions. — We have investigated the fundamental question whether the groundstate energy of hard-core bosons is lower than that of fermions on the same lattice. We have proved that the former is indeed lower than the latter, as naturally expected, if there is no frustration in the hopping. However, the inequality can be reversed in the presence of frustration, and we have demonstrated that it is actually the case in several concrete examples. There is a close connection of the present problem to many apparently unrelated concepts in quantum many-body problem, including Simon's universal diamagnetism of bosons, Lieb's optimal flux for fermions, statistical transmutation, flat-band ferromagnetism, and antiferromagnets near the saturation field. More details of our analysis will be presented in a separate publication [16].

We thank Ehud Altman for stimulating discussion which motivated us to study this problem. We also thank Tarun Grover for useful discussions. This work is supported in part by Grant-in-Aid for Scientific Research on Innovative Areas No. 20102008 from MEXT, Japan, and by US National Science Foundation Grant No. NSF PHY11-25915 during a visit of W. N. and M. O. to Kavli Institute for Theoretical Physics, UC Santa Barbara. H. K. was supported in part by Grand-in-Aid for Young Scientists (B) (Grant No. 23740298).

W. N. is supported by MEXT scholarship. Numerical calculations in this work were partially carried out using TITPACK ver.2 by H. Nishimori, and codes provided by ALPS project [22–24].

---

[1] S. D. Huber and E. Altman, Phys. Rev. B **82**, 184502 (2010).
[2] E. Altman (2010), private communications.
[3] F. Becca, L. Capriotti, S. Sorella, and A. Parola, Phys. Rev. B **62**, 15277 (2000).
[4] E. H. Lieb, T. Schultz, and D. J. Mattis, Ann. Phys. (N. Y.) **16**, 407 (1961).
[5] S. Katsura, Phys. Rev. **127**, 1508 (1962).
[6] P. Ginsparg, in *Fields, Strings and Critical Phenomena* (North-Holland, 1989).
[7] Y. Hatsugai, K. Ishibashi, and Y. Morita, Phys. Rev. Lett. **83**, 2246 (1999).
[8] G. W. Semenoff, Phys. Rev. Lett. **61**, 517 (1988).
[9] E. Fradkin, Phys. Rev. Lett. **63**, 322 (1989).
[10] Y. Saiga and M. Oshikawa, Phys. Rev. Lett. **96**, 036406 (2006).
[11] H. Tasaki, Phys. Rev. Lett. **69**, 1608 (1992).
[12] A. Mielke and H. Tasaki, Commun.Math.Phys. **158**, 341 (1993).
[13] M. E. Zhitomirsky and H. Tsunetsugu, Phys. Rev. B **70**, 100403 (2004).
[14] A. Mielke, Journal of Physics A: Mathematical and General **24**, 3311 (1991).
[15] D. L. Bergman, C. Wu, and L. Balents, Phys. Rev. B **78**, 125104 (2008).
[16] W. Nie, H. Katsura, and M. Oshikawa, in preparation.
[17] E. H. Lieb, Phys. Rev. Lett. **73**, 2158 (1994).
[18] B. Simon, Phys. Rev. Lett. **36**, 1083 (1976).
[19] P. W. Anderson, Phys. Rev. **83**, 1260 (1951).
[20] R. Tarrach and R. Valentí, Phys. Rev. B **41**, 9611 (1990).
[21] R. Valentí, P. J. Hirschfeld, and J. C. Anglés d'Auriac, Phys. Rev. B **44**, 3995 (1991).
[22] ALPS Project `http://alps.comp-phys.org/`.
[23] A. Albuquerque, F. Alet, P. Corboz, P. Dayal, A. Feiguin, S. Fuchs, L. Gamper, E. Gull, S. Gürtler, A. Honecker, et al., J. Magn. Magn. Mater. **310**, 1187 (2007).
[24] M. Troyer, in *Lecture Notes in Computer Science* (Springer, 1999), vol. 1732, p. 164.